

ЎЗБЕКИСТОН РЕСПУБЛИКАСИ  
ОЛИЙ ВА ЎРТА МАХСУС ТАЪЛИМ ВАЗИРЛИГИ

---

---

ФАРҒОНА ДАВЛАТ УНИВЕРСИТЕТИ

**FarDU.  
ILMIY  
XABARLAR-**

1995 йилдан нашр этилади  
Йилда 6 марта чиқади

3-2019

**НАУЧНЫЙ  
ВЕСТНИК.  
ФерГУ**

Издаётся с 1995 года  
Выходит 6 раз в год

Аниқ ва табиий фанлар

МАТЕМАТИКА

<b>У.Бекбаев, К.Муминов</b> Матрицали дифференциал тенгламалар системасини сиртлар учун Лоренц алмаштиришлари аниқлигида ечиш.....	5
<b>Э.Каримов, С.Кербал</b> Суб-диффузия ва тўлқин тенгламаларидан иборат аралаш типдаги тенглама учун трикоми типдаги масала.....	10

КИМЁ

<b>Н.Бозоров, В.Кудышкин</b> Метилакрилатнинг акрил кислотаси билан сополимеризацияси .....	15
<b>Ҳ.Исмоилова, Д.Бекчанов, Ш.Ҳасанов, М.Балтаева</b> Пластикат поливинилхлоридни полиэтиленполиамин билан модификациялаб олинган ионитга мис(II), никель (II) ва кобальт(II) ионларининг сорбцияси .....	19
<b>М.Ахмадалиев, И.Асқаров</b> Зарарсиз толалар асосида маҳсулот олиш канцероген асбест муаммоларининг ечими сифатида .....	22
<b>Б.Саттарова, И.Асқаров, Ш.Абдуллоев</b> Товуқ гўштини сертификатлашда унинг таркибидаги антиоксидантлар миқдорини аниқлаш.....	27

БИОЛОГИЯ, ҚИШЛОҚ ХЎЖАЛИГИ

<b>Р.Максудов</b> Фарғона вилоятида анорчиликни ривожлантириш ва ушбу соҳанинг истиқболлари .....	33
--	----

Ижтимоий-гуманитар фанлар

ИҚТИСОДИЁТ

<b>А.Гафуров, О.Гафуров</b> Агроиқтисодиётнинг муҳим моддий ва маънавий тимсоли (КФК 80 ёшда).....	36
---	----

ТАРИХ

<b>Б.Абдуллаев, З.Раҳманов, Н.Камбаров</b> Қўштепа-2 ёдгорлигидаги кўшк хандагининг тадқиқотлари .....	40
<b>У.Мирзалиев</b> Совет даврида Сирдарё вилоятидаги тарихий-демографик жараёнларнинг ўзига хос хусусиятлари .....	44
<b>И.Хўжахонов</b> Совет тарихшунослигида ўзбеклар миллий идентиклиги муаммосининг ўрганилиш жиҳатлари.....	48
<b>М.Хасанов, Юнпенг Танг, М.Хомиджонова</b> Шимолий Бақтриянинг Кушонлар даврига оид янги ёдгорликлари.....	54
<b>Р.Мамадалиев</b> Исмоил ака – темуршунос олим .....	59
<b>Ш.Охунжонова</b> XIX аср ўрталари - XX аср бошларида Фарғона водийси бозорлари тарихидан.....	64

ФАЛСАФА, СИЁСАТ

<b>Г.Ғаффарова</b> Мураккаб тизимларга оид илмий ғоялар .....	68
<b>М.Маматов</b> Тасаввуф таълимотининг ижтимоий-маданий детерминантлари .....	73
<b>Т.Абдуллаев, Б.Холматова</b> Инсон омилини фаоллаштириш масалалари .....	77
<b>О.Бойбуваева</b> Мамлакатимизда амалга оширилаётган диний-маърифий соҳадаги ислохотларнинг такомиллашуви .....	82

УДК: 517.956

СУБ-ДИФФУЗИЯ ВА ТЎЛҚИН ТЕНГЛАМАЛАРИДАН ИБОРАТ АРАЛАШ ТИПДАГИ  
ТЕНГЛАМА УЧУН ТРИКОМИ ТИПИДАГИ МАСАЛАTRICOMI TYPE PROBLEM FOR MIXED TYPE EQUATION WITH SUB-DIFFUSION AND WAVE  
EQUATIONЗАДАЧА ТИПА ТРИКОМИ ДЛЯ УРАВНЕНИЯ СМЕШАННОГО ТИПА С УРАВНЕНИЕМ СУБ-  
ДИФФУЗИИ И ВОЛНОВЫМ УРАВНЕНИЕМ

E.Karimov, S.Kerbal

**Аннотация**

Мақолада икки тартибли Хилфер ҳосилалари суб-диффузия ва классик тўлқин тенгламасида иборат аралаш типдаги тенглама учун Трикоми типдаги масала аралаш соҳада ўрганилган. Асосий қўлланилган усуллар интеграл тенгламалар ва энергия интеграллари усулларидир.

**Annotation**

In this paper, we have considered a Tricomi type problem for mixed type equation with Hilfer's double order derivative sub-diffusion equation and classical wave equation in a composite domain. Main methods of the investigation are a method of integral equations and energy integrals' method.

**Аннотация**

В статье рассмотрена задача типа Трикоми для уравнения смешанного типа с уравнением суб-диффузии и волновым уравнением. Основными методами являются метод интегральных уравнений и метод интегралов энергии.

**Таянч сўз ва иборалар:** Трикоми масаласи, икки тартибли Хилфер ҳосиласи, интеграл тенгламалар усули.

**Keywords and expressions:** Tricomi problem, double order Hilfer's derivative, method of integral equations

**Ключевые слова и выражения:** задача Трикоми, производное Хилфера с двумя порядками, метод интегральных уравнений.

**Introduction and formulation of a problem.**

Due to both practical [1] and theoretical importance [2], Fractional Calculus is developing intensively.


Solutions of differential equations with the most popular fractional derivatives such as the Riemann-Liouville, Caputo and Hilfer's derivatives were subject of investigations in [3], [4], [5]. Analyzing a huge amount of works devoted to the study of direct and inverse problems for partial differential equations (PDEs) with fractional order operators, we noted the only study closely related to the present topic. Several boundary value problems (BVPs) for mixed type equations with Riemann-Liouville fractional differential operator (FDO) were studied for unique solvability in works [6],[7]. BVP with integral form conjugation conditions for PDEs with both Riemann-Liouville and Caputo FDOs were subject of series of investigations [8], [9], [10]. In these works, authors used an explicit solution of certain BVP for fractional diffusion equation studied by A.Pskhu [5].

For the first time, generalized the Riemann-Liouville FDO, which is alternatively named as Hilfer FDO introduced by Hilfer [11]. The Cauchy and some BVPs for ODEs and PDEs with Hilfer FDO investigated by many authors, for instance [4], [12].

In this paper, we are aimed to study BVP with integral form conjugation condition in a mixed domain consisted of characteristic triangle and rectangle, for a mixed type PDE with diffusion equation involving Hilfer's double order FDO, which is introduced in [13]. In recent work [14], Tricomi type problem with integral conjugation condition for mixed type equation with Hilfer's derivative was investigated.

Consider the following mixed type equation

*E.Karimov – senior researcher of Uzbekistan Academy of Science Institute of Mathematics named after V.I Romanovski.  
S.Kerbal – professor of Sultan Qaboos University, Department of Mathematics, FarcDiff Research group, Muscat, Oman.*



$$0 = \begin{cases} D_{0t}^{(\alpha,\beta)\mu} u(x,t) - u_{xx}(x,t), & t > 0, \\ u_{tt}(x,t) - u_{xx}(x,t), & t < 0 \end{cases} \quad (1)$$

in a composite domain  $\Omega = \Omega_1 \cup \Omega_2 \cup AB$ . Here

$$D_{0t}^{(\alpha,\beta)\mu} f = I_{0t}^{\mu(1-\alpha)} \frac{d}{dt} I_{0t}^{(1-\mu)(1-\beta)} f \quad (0 < \alpha, \beta \leq 1; 0 \leq \mu \leq 1)$$

is the Hilfer's double-order fractional derivative of orders  $\alpha$  and  $\beta$  of type  $\mu$  [13],

$$I_{0t}^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t \frac{f(z) dz}{(t-z)^{1-\alpha}} \quad \text{is the Riemann-Liouville fractional integral of order } \alpha \text{ [2].}$$

**Problem.** To find a function  $u(x,t)$ , which is continuous in  $\bar{\Omega} \setminus AB$ , its Hilfer's double-order derivative is continuous in  $\Omega_1$  and it has continuous second order partial derivatives in  $\Omega_2$ , and it satisfies Eq. (1) in  $\Omega$  together with boundary conditions

$$u(0,t) = u(1,t) = 0, \quad 0 \leq t \leq T, \quad (2)$$

$$u(x/2, -x/2) = \psi(x), \quad 0 \leq x \leq 1, \quad (3)$$

conjugation conditions on  $AB$

$$\lim_{t \rightarrow +0} t^{(1-\mu)(1-\beta)} u(x,t) = u(x,-0), \quad 0 \leq x \leq 1, \quad (4)$$

$$\lim_{t \rightarrow +0} t^{1-\alpha} \left( t^{(1-\mu)(1-\beta)} u(x,t) \right)_t = u_t(x,-0), \quad 0 < x < 1. \quad (5)$$

Here  $\psi(x)$  is a given function such that  $\psi(0) = 0$ .

**1. Main functional relations.**

Introduce a notation

$$\tau_1(x) = \lim_{t \rightarrow +0} t^{(1-\mu)(1-\beta)} u(x, t), \quad 0 \leq x \leq 1. \tag{6}$$

Solution of (1) in  $\Omega_1$  satisfying condition (2) and (6) has a form [13]

$$u(x, t) = \int_0^1 \tau_1(\xi) M(x, \xi; t) d\xi, \tag{7}$$

where

$$M(x, \xi; t) = 2t^{(1-\mu)(\beta-1)} \Gamma(\alpha + \mu(1-\beta)) \sum_{k=1}^{\infty} E_{\beta+\mu(\alpha-\beta), \beta+\mu(1-\beta)} \left( -(k\pi)^2 t^{\beta+\mu(\alpha-\beta)} \right) \sin k\pi x \sin k\pi \xi, \tag{8}$$

$E_{\alpha, \beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)}$  is a two-parameter Mittag-Leffler function[2].

Doing the same as in work [14], we deduce

$$v_1(x) = \frac{(\beta + \mu(\alpha - \beta)) \Gamma(\alpha + \mu(1 - \beta))}{\Gamma(\beta + \mu(1 - \beta))} \tau_1''(x), \quad 0 < x < 1, \tag{9}$$

where  $v_1(x) = \lim_{t \rightarrow +0} t^{1-\alpha} \left( t^{(1-\mu)(1-\beta)} u(x, t) \right)_t$ ,  $0 < x < 1$ .

Solution of (1) in  $\Omega_2$  satisfying initial conditions

$$u(x, -0) = \tau_2(x), \quad 0 \leq x \leq 1, \quad u_t(x, -0) = v_2(x), \quad 0 < x < 1 \tag{10}$$

can be written by D'Halembert's formula

$$u(x, t) = \frac{1}{2} \left[ \tau_2(x-t) + \tau_2(x+t) + \int_{x-t}^{x+t} v_2(z) dz \right]. \tag{11}$$

Substituting (11) into (3) we deduce

$$v_2(x) = \tau_2'(x) - 2\psi'(x), \quad 0 < x < 1. \tag{12}$$

Relations (9) and (12) are main functional relations, which are essential in further.

## 2. A unique solvability of the problem.

Based on conjugation conditions (4), (5) from main functional relations (9) and (12) yields

$$\tau_1''(x) - A\tau_1'(x) = 2A\psi'(x), \quad 0 < x < 1, \tag{13}$$

where  $A = \frac{\Gamma(\beta + \mu(1 - \beta))}{(\beta + \mu(\alpha - \beta)) \Gamma(\alpha + \mu(1 - \beta))}$ . Boundary conditions in (2) yield

$$\tau_1(0) = \tau_1(1) = 0. \tag{14}$$

Problem (13)-(14) has an explicit solution in a form of

$$\tau_1(x) = 2 \int_0^1 \psi'(\xi) G(x, \xi) d\xi \tag{15}$$

with

$$G(x, \xi) = \frac{1}{e^{Ax} - e^{A(x-1)}} \begin{cases} (1 - e^{A\xi})(1 - e^{A(x-1)}), & 0 \leq \xi \leq x, \\ (1 - e^{A(\xi-1)})(1 - e^{Ax}), & x \leq \xi \leq 1. \end{cases} \quad (16)$$

Using representation (15), unknown function  $v_1(x)$  will be found by formula (9). Due to conditions (4), (5), other two unknown functions  $\tau_2(x)$ ,  $v_2(x)$  will be determined explicitly. The solution of the considered problem will be recovered by formulas (7) and (11) in domains  $\Omega_1$  and  $\Omega_2$ , respectively.

A uniqueness of the solution of the problem can be proved in a similar way as it is done in [14]. Namely, we multiply (9) by function  $\tau_1(x)$  and integrate along  $AB$ , then easily will get

$$A \int_0^1 \tau_1(x) v_1(x) dx + \int_0^1 (\tau_1'(x))^2 dx = 0. \quad (17)$$

Considering (4), (5) and (12) one can readily prove that  $\int_0^1 \tau_1(x) v_1(x) dx \geq 0$ . Since  $\beta + \mu(\alpha - \beta) > 0$ , then from (17) it follows that  $\tau_1(x) = 0$ . Further, considering solution of the first BVP for Eq.(1) in  $\Omega_1$ , we will get  $u(x, t) = 0$  in  $\Omega_1$ . Due to (4), one can easily deduce that  $u(x, t) = 0$  in  $\Omega$ .

Finally, we are able now to formulate our result as the following

**Theorem.** If  $\psi(x) \in C[0, 1] \cap C^1(0, 1)$ , then formulated problem has a unique solution represented as follows

$$u(x, t) = 2\theta(t) \int_0^1 \int_0^1 \psi'(\eta) G(\xi, \eta) M(x, \xi; t) d\xi d\eta + \\ + \theta(-t) \left[ \int_0^1 \psi'(\eta) [G(x-t, \eta) + G(x+t, \eta)] d\eta + \frac{1}{A} \int_{x-t}^{x+t} dz \int_0^1 \psi'(\eta) G_{zz}(z, \eta) d\eta \right],$$

where  $\theta(t) = 1$  for  $t \geq 0$  and  $\theta(t) = 0$  for  $t < 0$ .

#### References:

1. Uchaikin V. V. Fractional Derivatives for Physicists and Engineers. Vol. I: Background and Theory. Vol. II: Applications. Nonlinear Physical Science. Heidelberg: Springer and Higher Education Press, 2012.
2. Kilbas A. A., Srivastava H. M., Trujillo J. J. Theory and Applications of Fractional Differential Equations, volume 204. North-Holland Mathematics Studies. -Amsterdam: Elsevier, 2006.
3. Luchko Y., Gorenflo R. An operational method for solving fractional differential equations with the Caputo derivatives. Acta Mathematica Vietnamica, 24, 1999, pp.207- 233.
4. Sandev T., Metzler R., Tomovski Z. Fractional diffusion equation with a generalized Riemann-Liouville time fractional derivative. J. Phys. A: Math. Theor. 44, 2011, 255203 (21pp)
5. Pskhu A. V. Partial Differential Equations of Fractional Order (In Russian). Moscow: Nauka, 2005.
6. Gekkieva S. Kh. A boundary value problem for the generalized transfer equation with a fractional derivative in a semi-infinite domain (In Russian). Izv. Kabardino-Balkarsk. Nauchnogo Tsentra RAN 1(8), 2002, pp.6-8.
7. Kilbas A. A., Repin O. A. An analog of the Tricomi problem for a mixed type equation with a partial fractional derivative. Fract. Calc. Appl. Anal. 13(1), 2010, p.69-84

8. Berdyshev A. S., Cabada A., Karimov E. T. On a non-local boundary problem for a parabolic-hyperbolic equation involving Riemann-Liouville fractional differential operator. *Nonlinear Analysis*, 75, 2012, pp.3268-3273.
9. Agarwal P., Berdyshev A. S., Karimov E. T. Solvability of a non-local problem with integral transmitting condition for mixed type equation with Caputo fractional derivative. *Results in Mathematics*. 71(3), 2017, pp. 1235-1257
10. Karimov E. T., Berdyshev A. S., Rakhmatullaeva N. A. Unique solvability of a non-local problem for mixed-type equation with fractional derivative. *Mathematical Methods in the Applied Sciences*. 40(8), 2017, pp.2994-2999
11. Hilfer R. *Applications of Fractional Calculus in Physics*. Singapore: World Scientific, 2000.
12. Hilfer R., Luchko Y., Tomovski Z. Operational method for the solution of fractional differential equations with generalized Riemann-Liouville fractional derivatives. *Fract. Calc. Appl. Anal.* 12(3), 2009, pp.299-318
13. Bulavitsky V.M. Closed form of the solutions of some boundary-value problems for anomalous diffusion equation with Hilfer's generalized derivative. *Cybernetics and Systems Analysis*, Vol.30, No 4, 2014, 570-577.
14. Karimov E.T. Tricomi type boundary value problem with integral conjugation condition for a mixed type equation with Hilfer fractional operator. *Bulletin of the Institute of Mathematics*, No 1, 2019, 19-26.

*(Reviewer: A.K.O'rinov – doktor of physics and mathematics, professor).*