

O'ZBEKISTON RESPUBLIKASI  
OLIY VA O'RTA MAXSUS TA'LIM VAZIRLIGI  
FARG'ONA DAVLAT UNIVERSITETI

**FarDU.  
ILMIY  
XABARLAR-**

1995 yildan nashr etiladi  
Yilda 6 marta chiqadi

2-2022

**НАУЧНЫЙ  
ВЕСТНИК.  
ФерГУ**

Издаётся с 1995 года  
Выходит 6 раз в год

## FarDU. ILMIY XABARLAR – НАУЧНЫЙ ВЕСТНИК.ФЕРГУ

**Muassis:** Farg'ona davlat universiteti.

«FarDU. ILMIY XABARLAR – НАУЧНЫЙ ВЕСТНИК. ФерГУ» "Scientific journal of the Fergana State University" jurnali bir yilda olti marta elektron shaklda nashr etiladi.

Jurnal filologiya, kimyo hamda tarix fanlari bo'yicha O'zbekiston Respublikasi Oliy attestatsiya komissiyasining doktorlik dissertatsiyalari asosiy ilmiy natijalarini chop etish tavsiya etilgan ilmiy nashrlar ro'yxatiga kiritilgan.

Jurnaldan maqola ko'chirib bosilganda, manba ko'rsatilishi shart.

O'zbekiston Respublikasi Prezidenti Administratsiyasi huzuridagi Axborot va ommaviy kommunikatsiyalar agentligi tomonidan 2020 yil 2 sentabrda 1109 raqami bilan ro'yxatga olingan.

Muqova dizayni va original maket FarDU tahririy-nashriyot bo'limida tayyorlandi.

### Tahrir hay'ati

**Bosh muharrir**  
**Mas'ul muharrir**

SHERMUHAMMADOV B.SH.  
ZOKIROV I.I

FARMONOV Sh. (O'zbekiston)  
BEZGULOVA O.S. (Rossiya)  
RASHIDOVA S. (O'zbekiston)  
VALI SAVASH YYELEK (Turkiya)  
ZAYNOBIDDINOV S. (O'zbekiston)

JEHAN SHAHZADAH NAYYAR (Yaponiya)  
LEEDONG WOOK. (Janubiy Koreya)  
A'ZAMOV A. (O'zbekiston)  
KLAUS XAYNSGEN (Germaniya)  
BAXODIRXONOV K. (O'zbekiston)

G'ULOMOV S.S. (O'zbekiston)  
BERDISHEV A.S. (Qozog'iston)  
KARIMOV N.F. (O'zbekiston)  
CHESTMIR SHTUKA (Slovakiya)  
TOJIBOYEV K. (O'zbekiston)

### Tahririyat kengashi

QORABOYEV M. (O'zbekiston)  
OTAJONOV S. (O'zbekiston)  
O'RINOV A.Q. (O'zbekiston)  
KARIMOV E. (O'zbekiston)  
RASULOV R. (O'zbekiston)  
ONARQULOV K. (O'zbekiston)  
YULDASHEV G. (O'zbekiston)  
XOMIDOV G'. (O'zbekiston)  
DADAYEV S. (O'zbekiston)  
ASQAROV I. (O'zbekiston)  
IBRAGIMOV A. (O'zbekiston)  
ISAG'ALIYEV M. (O'zbekiston)  
TURDALIYEV A. (O'zbekiston)  
AXMADALIYEV Y. (O'zbekiston)  
YULDASHOV A. (O'zbekiston)  
XOLIQOV S. (O'zbekiston)  
MO'MINOV S. (O'zbekiston)  
MAMAJONOV A. (O'zbekiston)

ISKANDAROVA Sh. (O'zbekiston)  
SHUKUROV R. (O'zbekiston)  
YULDASHEVA D. (O'zbekiston)  
JO'RAYEV X. (O'zbekiston)  
KASIMOV A. (O'zbekiston)  
SABIRDINOV A. (O'zbekiston)  
XOSHIMOVA N. (O'zbekiston)  
G'OFUROV A. (O'zbekiston)  
ADHAMOV M. (O'zbekiston)  
XONKELDIYEV Sh. (O'zbekiston)  
EGAMBERDIYEVA T. (O'zbekiston)  
ISOMIDDINOV M. (O'zbekiston)  
USMONOV B. (O'zbekiston)  
ASHIROV A. (O'zbekiston)  
MAMATOV M. (O'zbekiston)  
SIDDIQOV I. (O'zbekiston)  
XAKIMOV N. (O'zbekiston)  
BARATOV M. (O'zbekiston)

**Muharrir:** Sheraliyeva J.

**Tahririyat manzili:**

150100, Farg'ona shahri, Murabbiylar ko'chasi, 19-uy.  
Tel.: (0373) 244-44-57. Mobil tel.: (+99891) 670-74-60  
Sayt: [www.fdu.uz](http://www.fdu.uz). Jurnal sayti

Bosishga ruxsat etildi:

Qog'oz bichimi: - 60×84 1/8

Bosma tabog'i:

Ofset bosma: Ofset qog'oz.

Adadi: 10 nusxa

Buyurtma №

FarDU nusxa ko'paytirish bo'limida chop etildi.

**Manzil:** 150100, Farg'ona sh., Murabbiylar ko'chasi, 19-uy.

**Farg'ona,**  
**2022.**

|  |     |
|--|-----|
| <b>J.Xudoyberdiyev, A.Reymov, R.Kurbaniyazov, SH.Namazov, O.Badalova, A.Seytnazarov</b><br>Qoraqalpog‘istonning jelvakli fosforit uni asosidagi faollashgan superfosfat .....                        | 248 |
| <b>M.Ahmedov, Z.Teshaboyev</b><br>“Hayot davomida o‘qish” tamoyili asosida innovatsion xarakterga ega<br>bo‘lgan “mavzu ishlanmasi” tayyorlash.....  | 255 |
| <b>KIMYO</b>   |     |
| <b>S.Samatov, A.Ikramov, O.Ziyadullayev, S.Abduraxmanova</b><br>Benzaldegid va uning xosilalarini fenilatsetilen ishtirokida enantioselektiv alkinillash jarayoni.....                               | 259 |
| <b>I.Asqarov, G‘.Madrahimov, M.Xojimatov</b><br><sup>3</sup> O-ferrotsenil benzoy kislotasini ayrim hosilalarining biologik faolligini o‘rganish .....   | 267 |
| <b>U.Mamatkulova, X.Isakov, I.Askarov</b><br>Sarimsoqpiyoz va po‘stining kimyoviy tarkibi, shifobaxshlik xususiyatlari.....  | 271 |
| <b>I.Askarov, Z.Nazirova</b><br>Qizil lavlagi tarkibidagi ayrim kimyoviy birikmalar va ularning ahamiyati .....  | 275 |
| <b>I.Asqarov, B.Nizomov</b><br>Yeryong‘oq va yong‘oq mevasining qiyosiy kimyoviy tarkibi va shifobaxsh xususiyatlari .....   | 279 |
| <b>I.Mamatova, I.Askarov</b><br>«Oltin vodiya» oziq-ovqat qo‘shilmasining giperglikemik xususiyatlari .....  | 283 |
| <b>I.Askarov, X.Isakov, SH.Turaxonov</b><br>Monometilolmochevinagallat efirlarini olish .....  | 286 |
| <b>BIOLOGIYA, QISHLOQ XO‘JALIGI</b>  |     |
| <b>A.Xusanov D.Kapizova, G.Zokirova, N.Oxunova</b><br>Farg‘ona vodiysi sharoitida ochiq urug‘li daraxt va butalarning so‘ruvchi fitofaglari (Iachnidae, diaspididae):<br>faunasi va ekologiyasi..... | 290 |
| <b>M.Nazarov, M.Ma‘murova, A.Xamidov M.Mirzaxalilov</b><br>Baliqchilik xo‘jaligi hovuzlarida fitoplankton tarkibi va o‘simlikxo‘r baliqlarni<br>yetishtirishda ularning o‘rni .....                  | 295 |
| <b>ILMIY AXBOROT</b>   |     |
| <b>Z.Xosilova</b><br>Oshiqcha tana vazni va uning aholi guruhlari (18-59 yoshlilar) orasida uchrash holati .....   | 299 |
| <b>S.Mamadaliyeva, M.Omonova, B.Saydaliyev</b><br>Mahalliy xomoshyodan adsorbentlarda parafinni chuqur tozalash uchun<br>kombinirlangan texnologiya .....  | 302 |
| <b>A.Xolikulov</b><br>Buxoro xonligining Rossiya bilan siyosiy aloqalari tarixidan.....  | 306 |
| <b>X.Jumaniyozov</b><br>Markaziy Osiyoning tabiiy-geografik, ijtimoiy-iqtisodiy, logistik imkoniyatlarining<br>geosiyosiy jarayonlarga ta‘siri.....  | 312 |
| <b>A.Hakimov</b><br>Sovet davri maktab o‘qituvchilarining kundalik hayoti .....  | 316 |
| <b>G‘.Israilov</b><br>Sakkokiy adabiy merosi o‘rganilishi manbalari xususida.....  | 321 |
| <b>S.Xoliqov</b><br>Milliy xavfsizlikni ta‘minlash jarayonida O‘zbekiston respublikasi Oliy Majlisi senati ishtirokining<br>tashkiliy va nazariy-huquqiy asoslari .....                              | 325 |
| <b>O.Axmadjonova</b><br>Badiiy asarda psixologik (ruhiy) tahlil printsiplari va usullari.....  | 330 |
| <b>D.Buzrukova</b><br>“Muhabbat” konseptining lingvomadaniy o‘ziga xosligi .....   | 334 |
| <b>K.Topvoldiyev</b><br>Lermontov M.Yu. asarlarida XIX asr kavkazi.....  | 338 |
| <b>S.Abduraxmonov, SH.Ibragimov</b><br>Ta‘lim tizimida baholashning asosiy mezonlari va uning ahamiyati .....  | 345 |
| <b>A.U.Choriyev, G.O Temirova,</b><br><b>Yetuk kimyogar, kamtarin olim</b> .....   | 349 |

## MONOMETILOLMOCHEVINAGALLAT EFIRLARINI OLISH

## ПОЛУЧЕНИЕ МОНОМЕТИЛОЛМОЧЕВИНАГАЛЛАТНЫХ ЭФИРОВ

## OBTAINING MONOMETHYLOLUREAGALLATE ETHERS

Askarov Ibrohim Rakhmonovich<sup>1</sup>, Isakov Khayatulla<sup>2</sup>,  
Turakhonov Shokhrukh Odiljonovich<sup>3</sup>

<sup>1</sup>Askarov Ibrohim Rakhmonovich

– Doctor of Chemistry Sciences, Professor of Andijan State University,

<sup>2</sup>Isakov Khayatulla

– Doctor of Technical Sciences, Professor of Andijan State University

<sup>3</sup>Turakhonov Shokhrukh Odiljonovich

– assistant of Andijan State University,

**Annotatsiya**

*Tadqiqotlar mochevina va formaldegidning o'zaro ta'siri natijasida mono-, di- va trimetilolmochevina hosil bo'lishini o'rgandi. Ushbu tadqiqotning maqsadi gal kislotasining mono-, di- va trimetilolmochevina bilan efirlarini hosil bo'lishini o'rganishdan iborat bo'ldi. Tadqiqotda mayda organik sintez, IK, mass-spektrometriya, erish nuqtasini o'lchash, yupqa qatlam, ustunli xromatografiya usullari qo'llanildi. Efirlarning tuzilishi shuni ko'rsatdiki, eterifikatsiya reaksiyasi gallyk kislotaning karboksil guruhi va metilol karbamidning gidroksil guruhi hisobiga boradi.*

**Аннотация**

*В исследованиях изучено образование моно-, ди- и триметилломочевин в результате взаимодействия мочевины и формальдегида. Целью данного исследования было изучение образования сложных эфиров галловой кислоты с моно-, ди- и триметилломочевин. В исследовании использованы методы тонкого органического синтеза, ИК-, масс-спектрометрия, измерение температуры плавления, тонкослойная, колоночная хроматография. Структура сложных эфиров показала, что реакция этерификации протекает за счет карбоксильной группы галловой кислоты и гидроксильной группы метилломочевин.*

**Abstract**

*The studies studied the formation of mono-, di- and trimethylol ureas as a result of the interaction of urea and formaldehyde. The purpose of this study was to study the formation of esters of gallic acid with mono-, di- and trimethylurea. The study used methods of fine organic synthesis, IR, mass spectrometry, melting point measurement, thin layer, column chromatography. The structure of esters showed that the esterification reaction proceeds due to the carboxyl group of gallic acid and the hydroxyl group of methylol ureas.*

**Kalit so'zlar:** mochevina, formal'degid, monometilolmochevina, ammiak, gall kislotasi, gallatlar, benzoic kislotasi, polifenollar va murakkab efirlar.


**Ключевые слова:** мочевины, формальдегид, монометилломочевина, аммиак, галловая кислота, галлаты, бензойная кислота, полифенолы и сложные эфиры.

**Keys words:** urea, formaldehyde, monomethylurea, ammonia, gallic acid, gallates, benzoic acid, polyphenols and esters.

**INTRODUCTION**

Methylolureas have been synthesized by different methods. Urea reacts with formaldehyde in an alkaline medium to form monomethylol urea (1) [1].

The reaction of urea with formaldehyde takes place in several stages. Their direction depends on temperature, pH value, concentration and time of reagent aggregation. At the first stage, methylol groups are formed. The presence of four reactive hydrogen atoms in urea indicates that it can combine with the same amount of formaldehyde.




Experiments have shown that the inclusion of each methylol group reduces the reactivity of the remaining hydrogen atoms in the amino and imine groups. The equilibrium constants of the reactions for the formation of mono-, di- and trimethylolureas are 0.045, 0.274 and 1.8 at 50 °C. The reactions for the formation of methylolurea are bimolecular, and the dissociation reaction is monomolecular. In these reactions,  $\text{H}_3\text{O}^+$ ,  $\text{OH}^-$ , acids, bases and salts act as catalysts. In the reaction of urea with formaldehyde, the rate of formation of monomethylolurea increases at 35 °C and pH 2.0-4.0 [1-3].

Gallic acid forms esters with alcohols. Esters of methyl, ethyl, propyl, and octyl alcohols with gallic acid were synthesized on a sulfuric acid catalyst.

Methyl gallate is oxidized to hexaoxidophenol. It is converted to ellagic acid by internal molecular transesterification in ammonia solution.

Ethanol is used in the fuel, polymer and cosmetic industries. It is not included in the list of approved food additives for use in the food industry in Russia and most countries of the world [4].

Since 1948, gallic acid propyl ester (1-propyl-3,4,5-trihydroxybenzoate) has been used as a food additive due to its antioxidant properties [5,6]. Gallic acid with benzoic acid forms benzoic gallate ester.



Gallic acid and its esters are used in the food industry as a food additive due to their antioxidant properties, in pharmaceuticals, medicine, in the production of paints and varnishes.

#### EXPERIMENTAL PART.

**Synthesis of monomethylurea (2).** 1200 g (20 mol) of urea was dissolved in 1200 ml of water. The aqueous solution was heated to 40 °C. The ammonia solution was added to 1615 ml of 34% formalin to a pH of 8-9 and the two solutions were mixed. When the mixture was kept at 10 °C for 4 h, a white precipitate formed. Then the solution was shaken and poured into a crystallizer. After 1 day, a white precipitate was isolated. M.p. = 100-101 °C.

**Synthesis of dimethylol urea (3).** 60 g (1 mol) of urea was dissolved in 60 ml of water. The aqueous solution was heated to 40 °C. The ammonia solution was added to 157.5 ml of 34% formalin to pH 8.3 and the two solutions were mixed. When the mixture was kept at 10 °C for 24 h, a white precipitate formed. Then the solution was shaken and poured into a crystallizer. After 1 day, a white precipitate was isolated. M.p. = 121-122 °C.

**Synthesis of trimethylol urea (4).** 60 g (1 mol) of urea was dissolved in 60 ml of water. The aqueous solution was heated to 40 °C. The ammonia solution was added to 290 ml of 34% formalin to a pH of 8.3 and the two solutions were mixed. When the mixture was kept at 10 °C for 24 h, a white precipitate formed. Then the solution was shaken and poured into a crystallizer. After 1 day, a white precipitate was isolated. M.p. = 127-128 °C.

Synthesis of gallic acid esters. At 0 °C, 2.5 g (12 mmol) DCC (dicyclohexylcarbodiimide) was slowly added to a solution of 1.7 g (10 mmol) gallic acid in 100 ml of diethyl ether. To the mixture was added a solution of 50 mmol of urea in 10 ml of diethyl ether, warmed to room temperature and stirred for 24 hours. The precipitate was separated from the reaction mixture, washed 3 times with 5% NaHCO<sub>3</sub>, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated in vacuo. The resulting mixture was separated by column chromatography (ethanol/chloroform 4:1).

Ureidomethylgallate (5) yield 46%. M.p.=155°C. *IR spectrum*. strong 580 cm<sup>-1</sup>, middle 765 cm<sup>-1</sup>, middle 880 cm<sup>-1</sup>, strong 940 cm<sup>-1</sup>, middle 990 cm<sup>-1</sup>, strong 1035 cm<sup>-1</sup>, strong 1090 cm<sup>-1</sup>, very strong 1110 cm<sup>-1</sup>, wide, strong 1130 cm<sup>-1</sup>, wide, strong 1190 cm<sup>-1</sup>, strong 1260 cm<sup>-1</sup>, strong 1570 cm<sup>-1</sup>, strong 1650 cm<sup>-1</sup>, powerless, shoulder 1670 cm<sup>-1</sup>, powerless 3280 cm<sup>-1</sup>, powerless 3355 cm<sup>-1</sup>, powerless 3370 cm<sup>-1</sup>, powerless 3720 cm<sup>-1</sup>, powerless 3770 cm<sup>-1</sup>, powerless 3830 cm<sup>-1</sup>, wide, powerless 3890-3900 cm<sup>-1</sup>. *Mass spectrum*: 243,0014 m/z (100 %), 242,0016 m/z (55 %), 244,0024 m/z (23 %).

(3-(hydroxymethyl)ureido) methylgallate (6) унуми 42 %. M.p.=167°C. *IR spectrum*. strong 584 cm<sup>-1</sup>, middle 697 cm<sup>-1</sup>, middle 875 cm<sup>-1</sup>, strong 935 cm<sup>-1</sup>, middle 986 cm<sup>-1</sup>, strong 1035 cm<sup>-1</sup>, strong 1090 cm<sup>-1</sup>, very strong 1110 cm<sup>-1</sup>, wide, strong 1125 cm<sup>-1</sup>, wide, strong 1180 cm<sup>-1</sup>, strong 1235 cm<sup>-1</sup>, strong 1567 cm<sup>-1</sup>, strong 1640 cm<sup>-1</sup>, powerless 1660 cm<sup>-1</sup>, powerless 3280 cm<sup>-1</sup>, powerless 3270 cm<sup>-1</sup>, powerless 3355 cm<sup>-1</sup>, powerless 3345 cm<sup>-1</sup>, powerless 3370 cm<sup>-1</sup>, powerless 3755 cm<sup>-1</sup>, wide, powerless 3890-3895 cm<sup>-1</sup>. *Mass spectrum*: 272,008 m/z (100 %), 273,001 m/z (55 %), 274,0012 m/z (23 %).

(3,3-bis(hydroxymethyl)ureido)methylgallate (7) унуми 37 %. M.p.=178°C. *IR spectrum*. middle 600 cm<sup>-1</sup>, middle 740, 795 cm<sup>-1</sup>, middle 900 cm<sup>-1</sup>, strong 925 cm<sup>-1</sup>, middle 860 cm<sup>-1</sup>, very strong 890 cm<sup>-1</sup>, strong 965 cm<sup>-1</sup>, very strong 925, 970 cm<sup>-1</sup>, wide, strong 1110, 1230 cm<sup>-1</sup>, wide, strong 1030, 1165, 1180 cm<sup>-1</sup>, strong 1250, 1335 cm<sup>-1</sup>, strong 1590 cm<sup>-1</sup>, strong 1690 cm<sup>-1</sup>, strong 1710 cm<sup>-1</sup>, powerless 3055-3135 cm<sup>-1</sup>, powerless 3155-3220 cm<sup>-1</sup>, powerless 3225-3275 cm<sup>-1</sup>, powerless, wide 3570 cm<sup>-1</sup>, wide, powerless 3570-3710 cm<sup>-1</sup>. *Mass spectrum*: 302,0014 m/z (100 %), 303,0016 m/z (55 %), 304,0024 m/z (23 %).

### THE DISCUSSION OF THE RESULTS.

Based on the biological activity of urea derivatives with formaldehyde and gallic acid, it can be assumed that the products of their interaction, in particular esters, also have high biological activity. For this purpose, the reactions of gallic acid esterification with monomethyl urea, dimethyl urea, trimethyl urea were studied.

The interaction of gallic acid and methylurea is an esterification reaction and is known to take place in a mild environment. The presence of oxidants in the environment leads to the oxidation of gallic acid, and in an alkaline environment - to the polycondensation of methylurea. Therefore, a soft water extractor in a neutral solvent will be required. In view of this, in the esterification reaction, a dichloromethane solvent was used in an argon atmosphere, dicyclohexylcarbodiimide as a water absorber, and dimethylaminopyridine as a protonating agent.

When the reaction mixture was tested by TLC, it was found that the formation of products began after 30 minutes. Due to the high solubility of monomethylurea in diethyl ether in comparison with other urea derivatives and the closeness of the R<sub>f</sub> values, a 4:1 mixture of ethanol and chloroform was used as an eluent in the separation of the reaction product from the primary monomethylurea by column chromatography. When separating the remaining esters, it was observed that a slightly easier separation occurs when using this system. It was found that as a reaction product of trimethylurea with gallic acid, two products were formed with very similar R<sub>f</sub> values (0.34 and 0.38). The first fraction (R<sub>f</sub> = 0.38) was formed in large quantities, and the yield of the second fraction was so low that it could not be separated in pure form. Therefore, the first fraction was used in subsequent analyzes.

The shift of the absorption line from 1760 cm<sup>-1</sup> to 1565-1570 cm<sup>-1</sup>, which is typical for stretching vibrations (ν) of the carbonyl group in the carboxyl of gallic acid in the IR spectra of the obtained substances, confirm the formation of an ester. In addition, a decrease in the peak characteristic of the valency variation (ν) of the OH group of gallic acid indicates that the hydroxyl group in the carboxyl is replaced by an alcohol residue. In the IR spectrum of 7 compounds, a

nonspecific decrease in the intensity of absorption lines 5 and 6 is observed. It can be argued that this is due to the esterification of a monosubstituted methylol group in trimethylolurea and the formation of intramolecular hydrogen bonds between di-substituted methylols in another amino group. This idea is also supported by the idea that the disubstituted methylols of the parent trimethylolurea are passive due to intramolecular hydrogen bonding.

In the mass spectra of the obtained substances, the high intensity of the peak belonging to the main molecular ion indicates the stability of the compounds.




Fig.1 Scheme of ureidogallates synthesis.

**CONCLUSIONS.** The fact that the synthesis of derivatives of methyl urea with gallic acid occurs under mild conditions, somewhat easier, leads to their further research. In particular, the study of the biological activity of the obtained esters will serve as the center of our future research.

#### References.

1. Асқаров И.Р., Хожиматов М.М., Исақов Х., Қирғизов. Метилломочевинанинг айрим ҳосилалари синтези//Илмий хабарнома. Андижон. -2014.-№4. -Б. 28-30. (Askarov IR, Khojimatov MM, Isakov H., Kirgizov. Synthesis of some derivatives of methyl urea // Scientific Bulletin. Andijan. -2014, №4, 28-30)
2. Асқаров И.Р., Исаев Ю.Т., Махсумов А.Г., Қирғизов Ш.М. Органик кимё.–Т.: F. Фулом, 2012. - 319-323б (Askarov I.R., Isaev Yu.T., Makhsumov AG, Kyrgyzov Sh.M. Organic chemistry. T.: G'. Gulom, 2012, 319-323).
3. Исақов Х., Усманов С., Горбунова В.В. ИК-спектроскопические исследования соединений тиомочевинны с формальдегидом // Узбекский химический журнал. -Тошкент.-1994. -№5. -С. 10-35. (Isafov X., Usmanov S., Gorbunova V.V. IR-spectroscopic studies of thiourea compounds with formaldehyde // Uzbek chemical journal, Tashkent, 1994, №5, 10-35)
4. Методы получения химических реактивов и препаратов. Москва, ИРЕА, Выпуск 15, 1967, стр. 187 (Methods of obtaining chemical reagents and drugs. Moscow, IREA, No 15, 1967, 187)
5. Final Report on the Amended Safety Assessment of Propyl Gallate // International Journal of Toxicology journal. - 2007. - Vol. 26, 3. - P. 89-118. (Заключительный отчет по дополненной оценке безопасности пропилгаллата // Международный журнал токсикологии. – 2007)
6. Jerker Widengren, Andriy Chmyrov, Christian Eggeling, Per-Åke Löfdahl, and Claus A. M. Seidel. Strategies to Improve Photostabilities in Ultrasensitive Fluorescence Spectroscopy // The Journal of Physical Chemistry journal A. - 2007. - Vol. 111, 3, 429-440. (Джеркер Виденгрэн, Андрей Чмыров, Кристиан Эггелинг, Пер-Оке Лёфдал и Клаус А. М. Зайдель. Стратегии улучшения фотостабильности в сверхчувствительной флуоресцентной спектроскопии // Журнал физической химии, журнал А. - 2007. - Том.)