

ЎЗБЕКИСТОН РЕСПУБЛИКАСИ  
ОЛИЙ ВА ЎРТА МАХСУС ТАЪЛИМ ВАЗИРЛИГИ

---

---

ФАРҒОНА ДАВЛАТ УНИВЕРСИТЕТИ

**FarDU.  
ILMIY  
XABARLAR-**

1995 йилдан нашр этилади  
Йилда 6 марта чиқади

— 2-2020 —

**НАУЧНЫЙ  
ВЕСТНИК.  
ФерГУ**

Издаётся с 1995 года  
Выходит 6 раз в год

Аниқ ва табиий фанлар

МАТЕМАТИКА

<b>Ш.Каримов, З.Комилова</b>	
Сингуляр коэффициентли тўртинчи тартибли битта тенглама учун Гурса масаласи.....	6
<b>К.Каримов</b>	
Яримчексиз параллелепипедда учта сингуляр коэффициентга эга бўлган эллиптик тенглама учун чегаравий масалалар .....	11

ФИЗИКА, ТЕХНИКА

<b>Р.Расулов, Б.Ахмедов, А.Абдухоликов, У.Раимжонова</b>	
Яримўтказгичларда ток ташувчилар эффектив Гамильтониани назарияси хусусида.....	24

КИМЁ

<b>М.Хожиматов, Ф.Абдугаппаров, И.Асқаров, Қ.Отахонов</b>	
М-ферроценилбензой кислотаси билан амигдалин реакциясини ўрганиш .....	28

БИОЛОГИЯ, ҚИШЛОҚ ХЎЖАЛИГИ

<b>Э.Исаков, Ш.Турдиев</b>	
Болалар орасида бирламчи ногиронлик структурасининг таҳлили.....	33
<b>Ф.Тухтасинов</b>	
Фарғона водийси жанубидаги сабзаёт экинларининг агробиоценозлари орасида тарқалган бегона ўтлардаги бўртма ва бошқа тур паразит фитонематодаларнинг тарқалиши .....	37

Ижтимоий-гуманитар фанлар

ФАЛСАФА, СИЁСАТ

<b>Т.Абдуллаев</b>	
Инсон фаолиятининг ижтимоийлашуви .....	42
<b>Г.Ғаффарова, Б.Қодиров</b>	
Ҳаракатлар стратегияси ва тизимли ёндашув .....	47
<b>М.Каримова</b>	
Инсоният онги ва қалбига қаратилган глобал таҳдидлар .....	52
<b>А.Музаффаров</b>	
Маданиятлараро мулоқот категорияси тадқиқига фалсафий-герменевтик ёндашув.....	56
<b>А.Қамбаров, Д.Тошалиев</b>	
Бузрук мақомидан Сарахбори бузрук шуъбасининг таҳлили масаласига доир .....	60

ТАРИХ

<b>Т.Эгамбердиева</b>	
1941-1945 йиллардаги Иккинчи жаҳон урушида ўзбек хотин-қизларининг маънавий жасорати.....	66
<b>Ҳ.Холиқулова</b>	
Ўзбекистон ногиронлар нодавлат нотижорат ташкилотлари: истиқболлари ва уларни ривожлантириш йўллари.....	71
<b>О.Ахмадов</b>	
Бухоро Халқ Совет Республикаси (БХСР)да таълим ва тарбияни замонавийлаштириш учун кураш .....	75

УДК:621. 315. 592

ЯРИМУТКАЗГИЧЛАРДА ТОК ТАШУВЧИЛАР ЭФФЕКТИВ ГАМИЛЬТониАНИ  
НАЗАРИЯСИ ХУСУСИДА

TO THE THEORY OF THE EFFECTIVE HAMILTONIAN OF CURRENT CARRIERS IN  
SEMICONDUCTORS

К ТЕОРИИ ЭФФЕКТИВНОГО ГАМИЛЬТониАНА НОСИТЕЛЕЙ ТОКА В  
ПОЛУПРОВОДНИКЕ

Р.Расулов, Б.Ахмедов, А.Абдухоликов, У.Раимжонова

**Аннотация**

Яримутказгич ўтказувчанлик, валент ва спин орбитал зоналарни ўз ичига олган Кейн модели ҳамда Латтинжер-Кон яқинлашишларида ток ташувчилар эффектив гамилтонианининг матрицавий элементлари таҳлил қилинган.

**Аннотация**

Рассчитаны матричные элементы эффективного гамилтониана носителей тока как в приближении Кейна, где рассматриваются зона проводимости, валентная зона, состоящая из подзон легких и тяжелых дырок и спин-отщепленная зона, так и в модели Латтинжера-Кона.

**Annotation**

The matrix elements of the effective Hamiltonian of current carriers are calculated as in the Kane approximation, where the conduction band, the valence band consisting of light and heavy hole subbands, and the spin-split band, as well as in the Luttinger-Kohn model, are considered.

**Таянч сўз ва иборалар:** матрицавий элемент, эффектив гамилтониан, ток ташувчилар, тўлқин функция.

**Ключевые слова и выражения:** матричный элемент, эффективный гамилтониан, носители тока, волновая функция.

**Keywords and expressions:** matrix element, effective Hamiltonian, current carriers, wave function.

It is known that many physical parameters of the crystalline potential depend on the band structure of the semiconductor [1, p.675; 2, p.435; 3, p.596; 4, p.1332; 5, p.560]. Moreover, usually in band theory it is believed that the crystalline periodic potential is always an even function of coordinates. However, in some cases, for example, in a semiconductor, where there is a heterojunction, the periodic potential of the crystal, along with the symmetric part, can have an asymmetric part.

This case requires a separate analysis of the matrix elements of the effective Hamiltonian of current carriers as in the Kane approximation, where the conduction band, the valence band consisting of light and heavy hole subbands, and the spin-split band, as well as in the Luttinger-Kohn model [6, p.484; 7, p.126]. Next, we consider the case when the extreme points of the zones are in the center of the Brillouin zone, i.e. at the point  $\vec{k} = 0$ , where  $\vec{k}$  is the wave vector of current carriers. In this case, the effective Hamiltonian can be represented as

$$H = H_0 + \frac{\hbar}{4m_0^2 c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} \quad (1)$$

and the corresponding (1) Schrödinger equation has the form

$$\left\{ H_0 + \frac{\hbar}{4m_0^2 c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} \right\} \psi_{n\vec{k}}(\vec{r}) = E_n(\vec{k}) \psi_{n\vec{k}}(\vec{r}), \quad (2)$$

where  $H_0 = \frac{p^2}{2m_0} + V(\vec{r})$  consists of kinetic and potential energy operators, the second term in (1) is the spin-orbit interaction operator,  $\vec{\sigma}$  is the vector of Pauli spin matrices with components:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (3)$$

**R.Rasulov** – professor of FSU, doctor of physics-mathematical sciences.

**B.Ahmedov** – FSU, doctorate student of physics department.

**A.Abdulolikov** – candidate for a master degree in radiophysics.

**U.Raimjonova** – a student of teaching methods of physics and astronomy.

whence for the spinors  $\uparrow \equiv \begin{pmatrix} 1 \\ 0 \end{pmatrix}$   $\downarrow \equiv \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  we have the following relations

$$\sigma_x \uparrow \downarrow \sigma_y \uparrow = i \downarrow \sigma_x \uparrow = \uparrow \sigma_x \downarrow = \uparrow \sigma_y \downarrow = -i \uparrow \sigma_x \downarrow = -\downarrow \quad (4)$$

If the solution (3) is sought in the form of the Bloch function  $\psi_{n\vec{k}}(\vec{r}) = e^{i\vec{k}\vec{r}} u_{n\vec{k}}(\vec{r})$ , then we obtain the equation for the Bloch amplitude  $u_{n\vec{k}}(\vec{r})$  as

$$\left\{ H_0 + \frac{\hbar}{m_0} \vec{k}\vec{p} + \frac{\hbar}{4m_0^2c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} + \frac{\hbar^2}{4m_0^2c^2} [\vec{\nabla}V \times \vec{k}] \cdot \vec{\sigma} \right\} u_{n\vec{k}}(\vec{r}) = E' u_{n\vec{k}}(\vec{r}) \quad (5)$$

where  $E' = E_n(\vec{k}) - \frac{\hbar^2 k^2}{2m_0}$ . The last term in (5) describes the spin-orbit interaction, which depends on the wave vector of current carriers. Thus, the effective Hamiltonian acting on the periodic function  $u_{n\vec{k}}(\vec{r})$  is expressed as:

$$H = H_0 + \frac{\hbar}{m_0} \vec{k}\vec{p} + \frac{\hbar^2}{4m_0^2c^2} [\vec{\nabla}V \times \vec{k}] \cdot \vec{\sigma} + \frac{\hbar}{4m_0^2c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} \quad (6)$$

Here  $H_1 = \frac{\hbar}{m_0} \vec{k}\vec{p}$  and  $H_2 = \frac{\hbar^2}{4m_0^2c^2} [\vec{\nabla}V \times \vec{k}] \cdot \vec{\sigma}$  appear due to the transition from the Bloch function to the function  $u_{n\vec{k}}(\vec{r})$ , the term  $H_3 = \frac{\hbar}{4m_0^2c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma}$  describes  $\vec{p}$  dependent in-orbit interaction. The Bloch amplitude  $u_{n\vec{k}}(\vec{r})$  for electrons in the conduction band can be represented as:  $|S \uparrow\rangle, |S \downarrow\rangle$ , and for holes in the valence band  $-|X \uparrow\rangle, |X \downarrow\rangle, |Y \uparrow\rangle, |Y \downarrow\rangle, |Z \uparrow\rangle, |Z \downarrow\rangle$  with the corresponding intrinsic energies  $E_s$  and  $E_p$ , which are defined as  $H_0|S\rangle = E_s|S\rangle, H_0|X\rangle = E_p|X\rangle, H_0|Y\rangle = E_p|Y\rangle, H_0|Z\rangle = E_p|Z\rangle$ , where [8, p. 688]

$$|S\rangle = \frac{1}{\sqrt{4\pi}} \cdot |Z\rangle = \sqrt{\frac{3}{4\pi}} \frac{z}{r}, |X \pm iY\rangle = \sqrt{\frac{3}{4\pi}} \frac{x \pm iy}{r}, \quad (7)$$

where it was considered that the wave function of the electrons in the conduction band is the wave function of the s-state, and for the valence band, the p-state of the hydrogen atom. Since the states in the conduction band are twofold degenerate along the spin, and in the valence band fourfold degenerate, therefore, the basic functions can be represented as:

$$|1\rangle = |S \downarrow\rangle, |2\rangle = |\frac{X-iY}{\sqrt{2}} \uparrow\rangle, |3\rangle = |Z \downarrow\rangle, |4\rangle = |-\frac{X+iY}{\sqrt{2}} \uparrow\rangle, \quad (8)$$

$$|\bar{1}\rangle = |S \uparrow\rangle, |\bar{2}\rangle = |-\frac{X+iY}{\sqrt{2}} \downarrow\rangle, |\bar{3}\rangle = |Z \uparrow\rangle, |\bar{4}\rangle = |\frac{X-iY}{\sqrt{2}} \downarrow\rangle. \quad (9)$$

First, we determine the diagonal matrix elements of the Hamiltonian (6) from the basis functions (8) and (9). This requires calculating the matrix elements of each term (6) separately, where in further calculations we take into account that  $\iiint_{-\infty}^{\infty} \frac{x^{2m+1} y^l z^\mu}{r^n} d\vec{r} = \iiint_{-\infty}^{\infty} \frac{x^m y^{2l+1} z^\mu}{r^n} d\vec{r} = \iiint_{-\infty}^{\infty} \frac{x^m y^l z^{2\mu+1}}{r^n} d\vec{r} = 0$ , where  $d\vec{r} = dx dy dz$ ,  $m, l, \mu$  are integers. Then the matrix elements of the operators

$$H_1 = \frac{\hbar}{m_0} \vec{k}\vec{p}, H_2 = \frac{\hbar^2}{4m_0^2c^2} [\vec{\nabla}V \times \vec{k}] \cdot \vec{\sigma}, H_3 = \frac{\hbar}{4m_0^2c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} \quad (10)$$

are defined with the following relations

$$\langle H_0 \rangle_{11} = \langle 1|H_0|1\rangle = \langle -iS \downarrow | H_0 | iS \downarrow \rangle = \langle S | E_s | S \rangle = E_s, \quad (11)$$

$$\langle H_1 \rangle_{11} = \langle 1|H_1|1\rangle = \langle -iS \downarrow | H_1 | iS \downarrow \rangle = \langle S | \frac{\hbar}{m_0} \vec{k}\vec{p} | S \rangle = 0, \quad (12)$$

$$\begin{aligned} \langle H_2 \rangle_{11} &= \langle 1|H_2|1\rangle = \langle -iS \downarrow | H_2 | iS \downarrow \rangle = \\ &= \langle -iS \downarrow | \frac{\hbar^2}{4m_0^2c^2} [\vec{\nabla}V \times \vec{k}] \cdot \vec{\sigma} | iS \downarrow \rangle = -\frac{\hbar^2}{16\pi m_0^2 c^2} J_1 \end{aligned} \quad (13)$$

where

$$J_1 = \iiint_{-\infty}^{\infty} \left\{ \frac{\partial V}{\partial x} k_y - \frac{\partial V}{\partial y} k_x \right\} dx dy dz \quad (14)$$

and take into account that  $\vec{p}|S\rangle = 0$  (since the function S is a constant value), as well as the conditions of orthonormal spinors

$$\sigma_x \uparrow \downarrow, \sigma_y \uparrow = i \downarrow, \sigma_x \uparrow = \uparrow, \sigma_x \downarrow = \uparrow, \sigma_y \downarrow = -i \uparrow, \sigma_z \downarrow = -\downarrow. \quad (15)$$

If we consider that the crystalline periodic potential consists of two: even and odd terms with respect to the coordinate inversion:  $V(\vec{r}) = V_{ass}(\vec{r}) + V_{sim}(\vec{r})$ , where  $V_{sim}(\vec{r})V(\vec{r}) = (-\vec{r})$ ,  $V_{ass}(\vec{r}) = -V_{ass}(-\vec{r})$ ,  $V_{ass}(\vec{r})$ , then it is easy to verify that the integral  $J_1$  has nonzero terms. Therefore, we analyze the following cases.

It follows from (14) that: a) if  $V(\vec{r})$  has an odd term with respect to z, then  $J_1 = 0$ ; b) if  $V(\vec{r})$  has an odd term with respect to x, then  $J_{11} \neq 0$ ; c) if  $V(\vec{r})$  has an odd term with respect to y, then  $J_{12} \neq 0$ ; e) if  $V(\vec{r})$  has an odd term with respect to x and y, then

$$J_1 \neq 0. \langle H_3 \rangle_{11} = \langle -iS \downarrow | H_3 | iS \downarrow \rangle = \langle -iS \downarrow | \frac{\hbar}{4m_0^2c^2} [\vec{\nabla}V \times \vec{p}] \cdot \vec{\sigma} | iS \downarrow \rangle =$$

$$= -\frac{\hbar}{4m_0^2 c^2} (S \left[ [\vec{\nabla} V \times \vec{p}]_z \right] S) = 0 \quad (16)$$

The diagonal matrix elements of the effective Hamiltonian are defined by the following expressions:

$$\begin{aligned} (H_0)_{22} &= \langle \frac{X+iY}{\sqrt{2}} | H_0 | \frac{X-iY}{\sqrt{2}} \rangle = E_p, \quad (17) \\ (H_2)_{22} &= \langle \frac{X+iY}{\sqrt{2}} \uparrow | H_2 | \frac{X-iY}{\sqrt{2}} \uparrow \rangle = J_{22}^{(1)} + J_{22}^{(2)} + J_{22}^{(3)} + J_{22}^{(4)}. \end{aligned}$$

where

$$\begin{aligned} J_{22}^{(1)} &= -\frac{\hbar^2}{4m_0^2 c^2} \frac{1}{i 4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \{x^2 y k_y + (xy^2 + xz^2) k_x\} dx dy dz, \\ J_{22}^{(2)} &= +\frac{\hbar^2}{4m_0^2 c^2} \frac{1}{i 4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \{(yx^2 + yz^2) k_y + xy^2 k_x\} dx dy dz, \\ J_{22}^{(3)} &= -\frac{\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \{(x^3 + xz^2) k_y + x^2 y k_x\} dx dy dz, \\ J_{22}^{(4)} &= -\frac{\hbar^2}{4m_0^2 c^2} \frac{1}{i 4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \{xy^2 k_y + (y^3 + yz^2) k_x\} dx dy dz, \end{aligned}$$

whence it is clear that: a)  $J_{22}^{(1)}$  consists of three terms, the first of which is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , and the second is different from zero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ ; b)  $J_{22}^{(2)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ ; c)  $J_{22}^{(3)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{sim}(z)$ ; d)  $J_{22}^{(4)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ .

$$(H_2)_{22} = \langle \frac{X+iY}{\sqrt{2}} \uparrow | H_2 | \frac{X-iY}{\sqrt{2}} \uparrow \rangle = \mathcal{R}_{22}^{(1)} + \mathcal{R}_{22}^{(2)} + \mathcal{R}_{22}^{(3)} + \mathcal{R}_{22}^{(4)}, \quad (18)$$

where

$$\begin{aligned} \mathcal{R}_{22}^{(1)} &= \frac{\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \{x^2 \frac{\partial V}{\partial x} k_y - x^2 \frac{\partial V}{\partial y} k_x\} dx dy dz, \\ \mathcal{R}_{22}^{(2)} &= \frac{(-i)\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \{xy \frac{\partial V}{\partial x} k_y - xy \frac{\partial V}{\partial y} k_x\} dx dy dz, \\ \mathcal{R}_{22}^{(3)} &= \frac{i\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \{xy \frac{\partial V}{\partial x} k_y - xy \frac{\partial V}{\partial y} k_x\} dx dy dz, \\ \mathcal{R}_{22}^{(4)} &= \frac{\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \{y^2 \frac{\partial V}{\partial x} k_y - y^2 \frac{\partial V}{\partial y} k_x\} dx dy dz. \end{aligned}$$

The last relations show that: a)  $\mathcal{R}_{22}^{(1)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ , and the third for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ ; b)  $\mathcal{R}_{22}^{(2)}$  consists of three terms, the first of which is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ , and the third with  $V(\vec{r}) = V_{asim}(x) + V_{asim}(y) + V_{sim}(z)$ ; c)  $\mathcal{R}_{22}^{(3)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , and the second and third terms are nonzero when  $V(\vec{r}) = V_{asim}(x) + V_{asim}(y) + V_{sim}(z)$ ; d)  $\mathcal{R}_{22}^{(4)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ .

$$\begin{aligned} (H_2)_{22} &= \frac{\hbar}{4m_0^2 c^2} \{ \mathfrak{I}_{22}^{(1)} + \mathfrak{I}_{22}^{(2)} + \mathfrak{I}_{22}^{(3)} + \mathfrak{I}_{22}^{(4)} \} \\ \mathfrak{I}_{22}^{(1)} &= i \frac{3\hbar}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \left\{ \frac{\partial V(\vec{r})}{\partial y} z x^2 - \frac{\partial V(\vec{r})}{\partial x} x^2 y \right\} dx dy dz, \\ \mathfrak{I}_{22}^{(2)} &= i \frac{3\hbar}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \left\{ \frac{\partial V(\vec{r})}{\partial y} y^2 z + (yx^2 + yz^2) \frac{\partial V(\vec{r})}{\partial z} \right\} dx dy dz, \\ \mathfrak{I}_{22}^{(3)} &= + \frac{3\hbar}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \left\{ xyz \frac{\partial V(\vec{r})}{\partial y} + (x^3 + xz^2) \frac{\partial V(\vec{r})}{\partial z} \right\} dx dy dz, \\ \mathfrak{I}_{22}^{(4)} &= -\frac{3\hbar}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^4} \left\{ xyz \frac{\partial V(\vec{r})}{\partial y} - xy^2 \frac{\partial V(\vec{r})}{\partial x} \right\} dx dy dz, \end{aligned}$$

Analyzing the last relations, we have that: a)  $\mathfrak{J}_{\mathbb{B}\mathbb{B}}^{(1)}$  is nonzero at  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{asim}(z)$ ; b)  $\mathfrak{J}_{\mathbb{B}\mathbb{B}}^{(2)}$  consists of three terms, the first of which is nonzero at  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{asim}(z)$ , and the second is nonzero at  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ , and the third with  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ ; c)  $\mathfrak{J}_{\mathbb{B}\mathbb{B}}^{(3)}$  consists of three terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{asim}(z)$ , and the second and third terms are nonzero when  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ ; d)  $\mathfrak{J}_{\mathbb{B}\mathbb{B}}^{(4)}$  consists of two terms, the first of which is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{asim}(z)$ , and the second is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ .

Below are expressions for the matrix element of each member of the Hamiltonian. In particular,  $(H_0)_{\mathbb{B}\mathbb{B}} = \langle Z \downarrow | H_0 | Z \downarrow \rangle = \langle Z \downarrow | E_p | Z \downarrow \rangle = E_p$  and does not depend on the parity of the crystal potential relative to the coordinates;

$$(H_1)_{\mathbb{B}\mathbb{B}} = \langle Z \downarrow | H_1 | Z \downarrow \rangle = \langle Z \downarrow | \frac{\hbar}{m_0} \vec{k} \vec{p} | Z \downarrow \rangle = \langle \downarrow \downarrow \rangle \langle Z \downarrow | \frac{\hbar}{m_0} \vec{k} \vec{p} | Z \downarrow \rangle = \\ = i \frac{\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \left\{ k_x(-xz^2) + k_y(-yz^2) + k_z z \frac{y^2 + x^2}{r^2} \right\} dx dy dz,$$

whence we have that the first term of the last integral is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , the third is when  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ .

$$(H_2)_{\mathbb{B}\mathbb{B}} = \langle Z \downarrow | H_2 | Z \downarrow \rangle = \langle Z \downarrow | \frac{\hbar}{m_0} \vec{k} \vec{p} | Z \downarrow \rangle = \langle \downarrow \downarrow \rangle \langle Z \downarrow | \frac{\hbar}{m_0} \vec{k} \vec{p} | Z \downarrow \rangle = \\ = i \frac{\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \left\{ k_x(-xz^2) + k_y(-yz^2) + k_z z \frac{y^2 + x^2}{r^2} \right\} dx dy dz.$$

It can be seen from the last relations that the first term of the matrix element  $(H_2)_{\mathbb{B}\mathbb{B}}$  is nonzero for  $V(\vec{r}) = V_{asim}(x) + V_{sim}(y) + V_{sim}(z)$ , and the second for  $V(\vec{r}) = V_{sim}(x) + V_{asim}(y) + V_{sim}(z)$ , the third is when  $V(\vec{r}) = V_{sim}(x) + V_{sim}(y) + V_{asim}(z)$ .

$$(H_3)_{\mathbb{B}\mathbb{B}} = \langle Z \downarrow | H_3 | Z \downarrow \rangle = \langle Z \downarrow | \frac{\hbar^2}{4m_0^2 c^2} [\vec{p} \nabla \times \vec{p}] \cdot \vec{\sigma} | Z \downarrow \rangle = \\ = - \frac{i\hbar^2}{4m_0^2 c^2} \frac{3}{4\pi} \iiint_{-\infty}^{\infty} \frac{1}{r^2} \left\{ z^2 \frac{\partial v(\vec{r})}{\partial x} y - xz^2 \frac{\partial v(\vec{r})}{\partial y} \right\} dx dy dz \text{ and this matrix element is nonzero at } \\ V(\vec{r}) = V_{asim}(x) + V_{asim}(y) + V_{sim}(z).$$

Thus, it was shown that, when the asymmetric part of the crystalline potential in semiconductors is taken into account, additional terms are obtained in the matrix elements of the effective Hamiltonian.

If we assume that the crystalline potential does not have an asymmetric part, then all the expressions obtained above and related to  $V_{asim}(x, y, z)$  turn to zero automatically.

#### Literature

1. Charles Kittel. Introduction to Solid State Physics. John Wiley and Sons, Inc. All. Rights reserved. 2005.
2. Ziman J.M. Principles of the Theory of Solids. Cambridge University Press, 1972.
3. Mohammad Abdul Wahab. Solid State Physics: Structure and Properties of Materials Alpha Science International, -2005. -596 p. 1842652184, 9781842652183.
4. Neil W. Ashcroft, Mermin Ashcroft, Dan Wei, N.David Mermin. Solid State Physics. CENGAGE Learning Asia, -2016.
5. Cardona Yu Peter, Cardona Manuel. Fundamentals of Semiconductor Physics. Per. from English I.I. Reshina. Ed. B.P. Zakharcheni. - 3rd ed., Rev. and add. -M.: Fizmatlit, -2002. -560 s. <http://www.twirpx.com/file/221809/>.
6. Bir G. L., Pikus G. E. Symmetry and Strain-induced Effects in Semiconductors. Wiley, 1974 - 484 p. ISBN 0470073217, 9780470073216.
7. Ivchenko E.L., Rasulov R.Ya. Symmetry and real band structure of half-conductors. -T.: - Fan, 1989.
8. [8] Landau L.D., Lifshitz E.M. Quantum Mechanics. 3rd edition. Non-Relativistic Theory. Pergamon. -1977.